



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

MECHANICS.

237. Proposed by C. N. SCHMALL, 604 East 5th Street, New York.

In a naval action an officer observes that in the case of two guns firing, at elevations α and β , respectively, the projectiles of the former fall a feet short of the target while those of the latter land b feet beyond. The initial velocity being the same in both cases, prove that the *true* elevation is

$$\frac{1}{2}\sin^{-1}\left[\frac{a\sin 2\beta + b\sin 2\alpha}{a+b}\right].$$

(Suggested by problem 29, page 219, Jeans' *Theoretical Mechanics*.)

Solution by PROFESSOR F. L. GRIFFIN, Williams College.

Let R =horizontal distance to target, V =initial velocity, g =the gravitational constant, and ϕ =elevation of gun to make the range equal to R .

Then the standard formula for the horizontal range gives for the elevations of ϕ , α , and β :

$$\begin{aligned}(1) \quad & V\sin 2\phi = gR, \\(2) \quad & V\sin 2\alpha = g(R-a), \\(3) \quad & V\sin 2\beta = g(R+b).\end{aligned}$$

Multiplying (3) by α , (2) by b , and adding, we obtain

$$V(a\sin 2\beta + b\sin 2\alpha) = gR(a+b) = V\sin 2\phi(a+b),$$

$$\text{whence } \phi = \frac{1}{2}\sin^{-1}\left[\frac{a\sin 2\beta + b\sin 2\alpha}{a+b}\right].$$

Also solved by H. C. Feemster, J. Scheffer, G. B. M. Zeer, S. G. Barton, and J. E. Sanders.

238. Proposed by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa.

Find the position of the center of pressure of a semi-elliptical area completely immersed in water, the bounding major-axis being inclined to the horizon at an angle β , and having one extremity in the surface of the water.

Solution by the PROPOSER.

If the area is not in the same vertical plane as the major axis, suppose it is inclined at an angle α . The ellipse projects into an ellipse in a vertical plane having the semi-axes a and $b\cos \alpha$.

Then from 229, pp, 189-190, Vol. XVI, No. 11, we get

$$= \frac{a(16b\cos \alpha \cos \beta + 15a\pi \sin \beta)}{4(4b\cos \alpha \cos \beta + 3\pi a\sin \beta)},$$

$$\frac{-}{y} = \frac{b \cos \alpha (16 a \sin \beta + 3 \pi b \cos \alpha \cos \beta)}{4(4b \cos \alpha \cos \beta + 3 \pi a \sin \beta)}$$

The distance of the center of pressure below the surface of the water is

$$\frac{b \cos \alpha \cos \beta (16 a \sin \beta + 3 \pi b \cos \alpha \cos \beta) + a \sin \beta (16 b \cos \alpha \cos \beta + 15 a \pi \sin \beta)}{4(4b \cos \alpha \cos \beta + 3 \pi a \sin \beta)}$$

$$= \frac{32 a b \sin \beta \cos \beta \cos \alpha + 3 \pi b^2 \cos^2 \alpha \cos^2 \beta + 15 \pi a^2 \sin^2 \beta}{4(4b \cos \alpha \cos \beta + 3 \pi a \sin \beta)}.$$

Also solved by S. G. Barton.

239. Proposed by J. G. ROSE, B. A. (Oxon), Mt. Angel College, Oregon.

A uniform bar of length $2a$ is placed in a sloping position, its lower end on the ground (coefficient of friction being μ), its upper end in the air, the bar being supported by a rough fixed peg (coefficient of friction μ'), against which it rests. If h is the height of the peg from the ground, and if θ be the angle the bar makes with the horizon, when on the point of slipping, prove that θ is to be found from the equation

$$\sin \theta \cos \theta [(\mu - \mu') \cos \theta + \sin \theta (1 + \mu \mu')] = \mu h/a.$$

Solution by S. G. BARTON, Ph. D., Clarkson School of Technology, Potsdam, N. Y.

Let R and R' be the pressures on the ground and peg, respectively, and W the weight. Then resolving the forces vertically and horizontally, and taking moments about the point on the ground, we have the three equations:

$$(1) \quad W = R + R' (\cos \theta + \mu' \sin \theta),$$

$$(2) \quad \mu R + R' (\mu' \cos \theta - \sin \theta) = 0,$$

$$(3) \quad R' h \csc \theta = W a \cos \theta, \text{ or } W = \frac{R' h}{a \sin \theta \cos \theta}.$$

Substitute this value of W in (1), multiply the equation by μ and subtract (3) to eliminate R' , and we have

$$\mu \cos \theta + \mu \mu' \sin \theta - \frac{\mu h}{a \sin \theta \cos \theta} - \mu' \cos \theta + \sin \theta = 0.$$

$$\text{Whence, } \sin \theta \cos \theta [(\mu - \mu') \cos \theta + \sin \theta (1 + \mu \mu')] = \mu (h/a).$$

Also solved by G. B. M. Zerr and J. Scheffer.